

Efficacy of slow release N-fertilizers for rice and their residual effect on wheat

M. K. SINGH, U. N. VERMA, S. K. PAL AND R. THAKUR

Department of Agronomy, Birsa Agricultural University, Ranchi 834 006

Received: July 1994

ABSTRACT

A field experiment was conducted during 1989-91 on sandy clay loam soil at Ranchi to study the efficiency of forms of urea viz. prilled (PU), lac coated (LCU), rock phosphate coated (RCU), karanj cake coated (KCU) and neem cake coated (NCU) at varying N rates (40, 80, 120 kg/ha) on the productivity of rice and their residual effect on succeeding wheat. LCU had markedly higher grain yield of rice than other forms of urea. Each incremental N level recorded higher grain yield over its preceding level. However, nitrogen use efficiency decreased with increasing N levels. Application of LCU and NCU had higher soil available N after rice as compared with other forms of urea. This was reflected on the productivity of succeeding wheat. Rice fertilized with 120 kg N/ha also recorded the maximum grain yield of succeeding wheat.

Key words: Slow release N fertilizers, Rice, Residual effect, Wheat

The recovery of fertilizer N in rice does not exceed 50% as considerable amounts are lost through leaching, volatilization, run-off and denitrification. These losses can be minimised by nitrification inhibitors viz. AM and N-Serve (Prasad and Power, 1995) and coating of urea with nitrification retarders viz. neem cake, karanj cake and lac (Chakravarty, 1979). Since nitrification inhibitors are costly and limited in availability, the coating of urea with indigenous materials is of paramount importance, as they release nitrogen slowly to meet the demand of rice crop throughout the growing period and also leave significant residual effect on the succeeding crop in sequence. This study was undertaken to

determine the efficacy of urea coated with various indigenous materials at varying N rates on the productivity of rice and their residual effect on succeeding wheat.

MATERIALS AND METHODS

Field experiments were conducted at Birsa Agricultural University Farm, Ranchi in 1989-90 and 1990-91 on a sandy clay loam soil. The soil was low in available N (240 kg/ha), P (10 kg/ha) and K (140 kg/ha) with pH 6.5. Sixteen treatments consisting of five forms of urea viz. prilled (PU), lac coated (LCU), rock phosphate coated (RCU), karanj cake coated (KCU) and neem cake coated (NCU) with three levels of N (40, 80 and 120 kg/ha) including one control

were tested in split plot design keeping N sources and control in main plot and their rates in subplot with three replications. Full dose of N was applied at transplanting as per treatment except the PU which was applied in three splits (50% at transplanting, 25% at active tillering and 25% at panicle initiation). A basal dose of 60 kg P_2O_5 and 40 kg K_2O /ha was also applied. The rice variety Sita was transplanted on 8 and 17 July and harvested on 25 October and 2 November in 1989 and 1990, respectively. The succeeding wheat Sonalika was sown on 15 and 22 November in 1989 and 1990, respectively with 50% recommended fertilizers (50 : 25 : 12.5 kg. N : P_2O_5 : K_2O /

ha) and harvested on 25 and 29 March in 1990 and 1991, respectively.

RESULTS AND DISCUSSION

Yield of rice

Sources of N significantly affected the grain yield of rice during both the years (Table 1). Lac coated urea produced 24% more mean grain yield than PU (27.2 q/ha) and exhibited its superiority over rest of the N sources tested. NCU and KCU were being on par produced 14.2 and 12.7% more grain yield than PU. In terms of N use efficiency, LCU was most efficient followed by NCU and KCU as their magnitude of increase was 69, 40 and 36% compared with PU

Table 1. Effect of sources and levels of nitrogen on grain yield and nitrogen use efficiency of rice and their residual effect on succeeding wheat and soil available N after rice

Treatment	Grain yield (q/ha)						Mean NUE (kg grain/ kg N)		Soil available N after rice (kg/ha)	
	Rice			Wheat			Rice	Wheat	1989	1990
	1989	1990	Pooled	1989- 90	1990- 91	Pooled				
<i>Source of N</i>										
PU	26.2	28.1	27.2	16.5	15.4	15.9	11.9	4.1	238	230
LCU	30.9	36.6	33.7	22.6	21.8	22.2	20.1	12.0	285	274
RCU	26.7	28.5	27.6	17.1	16.1	16.6	12.4	4.9	247	235
KCU	29.0	32.3	30.6	21.0	20.6	20.8	16.2	10.2	265	260
NCU	29.2	32.8	31.0	21.9	21.2	21.5	16.7	11.1	280	271
CD (P = 0.05)	1.6	3.3	2.2	1.2	1.1	1.2			13	10
<i>N (kg/ha)</i>										
40	24.9	27.2	26.0	16.2	14.9	15.6	20.8	7.3	235	226
80	28.7	32.0	30.3	19.9	19.3	19.6	15.8	8.6	268	256
120	31.7	35.8	33.8	23.4	22.8	23.1	13.4	8.7	286	280
CD (P = 0.05)	1.8	2.6	2.4	2.0	1.9	2.2			16	12
<i>Control vs. rest</i>										
Control	16.5	18.8	17.7	12.8	12.5	12.7			220	212
Rest	28.4	31.7	30.0	19.8	19.0	19.4			263	254
CD (P = 0.05)	3.6	5.4	5.0	4.2	3.9	4.4			35	27

respectively.

Similarly levels of N positively affected the grain yield of rice as each incremental level produced higher grain yield over its preceding level. Mean grain yield (33.8 q/ha) was maximum at 120 kg N/ha. However, nitrogen use efficacy decreased with N rates as it was 20.8, 15.8 and 13.4 kg grain/kg N at 40, 80 and 120 kg applied N/ha respectively. This confirms the findings of Panigrahi and Dixit (1991).

Residual effect on wheat

Sources and levels of N applied to rice exhibited significant residual effect on the productivity of succeeding wheat crop. Application of LCU and NCU in rice produced more grain yield of succeeding wheat than PU because of greater availability of N after rice harvest (Table 1). Further NCU and KCU had similar carry-over effect on the productivity of wheat.

Increasing levels of N to rice exhibited increased carry-over effect and productivity of succeeding wheat was maximum (23.1 q/ha) after rice fertilized with 120 kg N/ha. Residual nitrogen utilization efficiency by wheat crop followed the trend of rice crop for different sources. However, the trend was reverse for N rate applied to rice indicating that residual nitrogen utilization efficiency by wheat increases with increasing rate of N applied to rice crop.

Soil available N after rice

Sources and levels of N applied to rice significantly affected the soil available N after crop harvest (Table 1). Lac and neem coated urea treated plots had significantly higher soil available N after rice crop than the plots receiving PU and RCU. The plots receiving KCU had intermediate value of soil available N.

Increasing N levels to rice crop increased the soil available N after rice but the magnitude of increase decreased with increasing N. Soil available N in control plot after rice was reduced by 10% from its initial value (240 kg/ha) indicating the significance of nitrogen application for sustaining soil fertility and productivity of the crop.

REFERENCES

- Chakravarty, S. P. 1979. Effect of application of urea and ammonium sulphate blended with neem cake and compost on N transformation in soil and yield and nutrition of rice. *Journal of Indian Society of Soil Science* 27 (4) : 449-451.
- Panigrahi, B. K. and Dixit, L. 1991. Relative efficiency of slow release nitrogenous fertilizers in rainfed rice (*Oryza sativa*). *Indian Journal of Agronomy* 36 (3) : 401-403.
- Prasad, R. and Power J. F. 1995. Nitrification inhibitors for agricultural health and the environment. *Advances in Agronomy* 54 : 233-281.